

- (54) Title
AMINO ACID DERIVATIVES
- (51)^a International Patent Classification(s)
 C07K 007/06 A61K 037/64 C07K 001/02 C07K 005/06
 C07K 005/08 C07K 005/10
- (21) Application No. : 12617/88 (22) Application Date : 01.03.88
- (30) Priority Data
- (31) Number (32) Date (33) Country
 3707879 12.03.87 DE FEDERAL REPUBLIC OF GERMANY
 3721855 02.07.87 DE FEDERAL REPUBLIC OF GERMANY
- (43) Publication Date : 15.09.88
- (44) Publication Date of Accepted Application : 19.09.91
- (71) Applicant(s)
MERCK PATENT GESELLSCHAFT MIT BESCHRANKTER HAFTUNG
- (72) Inventor(s)
JOACHIM GANTE; PETER RADDATZ; JOHANNES SOMBROEK; CLAUS- J. SCHMITGES; KLAUS OTTO MINCK
- (74) Attorney or Agent
ARTHUR S CAVE & CO , GPO Box 3876, SYDNEY NSW 2001
- (56) Prior Art Documents
 AU 22982/88 C07K 5/06 5/08 5/10
 AU 79823/87 C07K 5/06 5/08 5/10
 AU 76222/87 C07K 5/06 5/08 5/10
- (57) Claim

1. Amino acid derivatives of the formula I




wherein

- X is H , $R^1-O-C_mH_{2m}-CO-$, $R^1-C_mH_{2m}-O-CO-$,
 $R^1-C_mH_{2m}-CO-$, R^1-SO_2- or
 $(R^1-C_mH_{2m}-(T)_x-C_rH_{2r})-L(R^7-C_pH_{2p})-C_tH_{2t}-CO-$,
- Z is 0 to 4 amino acid radicals attached to one another by a peptide linkage and selected from the group consisting of Abu, Ada, Ala, BAla, Arg, Asn, Asp, Bia, Cal, Dab, Gln, Glu, Gly, His, N(im)-alkyl-His, Ile, Leu, tert.-Leu, Lys, Met, α Nal, BNal, Nbg, Nle, Orn, Phe, Pro, Ser, Thr, Tic, Trp, Tyr and Val,
- E is 0 to 2 amino acid radicals attached to one another by a peptide linkage and selected from the group consisting of Abu, Ala, Cal, His, Ile, Leu, Met, Nle, Phe, Trp, Tyr and Val,
- D is $-CH_2-CHOH-CH_2OH$, $-C_2H_2-SO_2-R^{14}$, a phenyl-

BEST AVAILABLE COPY

dyl-C_yH_{2y}- radical which is substituted by one or two R¹⁴-SO₂- groups or an R¹⁴-CO- group or an (R¹⁴)₂-PO- group and, if desired additionally by an Hal atom, or D is



R¹, R³, R⁷ and R⁸ are each H, A, Ar, Ar-alkyl, Het,

Het-alkyl or cycloalkyl having 3-7 C atoms, cycloalkylalkyl having 4-11 C atoms, bicycloalkyl or tricycloalkyl having in each case 7-14 C atoms or bicycloalkylalkyl or tricycloalkylalkyl having in each case 8-18 C atoms, each of which is unsubstituted or monosubstituted or polysubstituted by A, AO and/or Hal,

R², R⁵ and R⁶ are each H or A,

R⁴ is (H, OH), (H, NH₂) or =O,

R⁹ is H, NH₂, NHA or NA₂,

R¹⁰, R¹¹, R¹² and R¹³ are each H, Hal, OH, OA, NH₂, SH, SA, SO₂NH₂, CF₃, CN, COOH or COOA,

R¹⁴ is OH, OA, NH₂, NHA, NA₂, NHCycloalkyl having 3-7 C atoms, N(cycloalkyl)₂ having 6-14 C atoms, pyrrolidino, piperidino, hexahydroazepino, morpholino, thiomorpholino, piperazino, N-A-piperazino, NHA₂ or NHHet,

L is CH or N,

T is O, S, NH or NA,

n is 1 or 2,

m, p, r and t are each 0, 1, 2, 3, 4 or 5,

x is 0 or 1,

y is 0, 1 or 2,

z is 2, 3, 4, 5 or 6,

Ar is phenyl which is unsubstituted or mono-substituted

and/or NH₂ or unsubstituted naphthyl,
Het is a saturated or unsaturated 5-membered or 6-membered heterocyclic radical which has 1-4 N, O and/or S atoms, which can be condensed with a benzene ring and/or can be monosubstituted or polysubstituted by A, AO, Hal, CF₃, HO, O₂N, carbonyl oxygen, H₂N, HAN, A₂N, AcNH, AS, ASO, ASO₂, HOOC, AOC, CN, H₂NCO, H₂NSO₂, ASO₂NH, Ar or Ar-alkenyl, hydroxyalkyl and/or aminoalkyl having in each case 1-8 C atoms, and/or in which the N and/or S heteroatoms can also be oxidized,
Hal is F, Cl, Br or I,

Ac is A-CO-, Ar-CO- or A-NH-CO-,
alkyl- is an alkylene group having 1-4 C atoms and
A is alkyl having 1-8 C atoms,
and wherein it is also possible for one or more -NA-CO-groups to replace one or more -NH-CO- groups, and salts thereof.

AUSTRALIA

PATENTS ACT 1952

COMPLETE SPECIFICATION

(ORIGINAL)

FOR OFFICE USE

Application Number:
Lodged:

Complete Specification Lodged:
Accepted:
Published:

Priority:
Related Art:

TO BE COMPLETED BY APPLICANT

Name of Applicant: MERCK PATENT Gesellschaft mit
beschränkter Haftung

Address of Applicant: Frankfurter Strasse 250,
D-6100 Darmstadt, Federal
Republic of Germany

Actual Inventor:

1. Dr Joachim GANTE
2. Dr Peter RADDATZ
3. Dr Johannes SOMBROEK
4. Dr Claus- J SCHMITGES
5. Dr Klaus Otto MINCK

Address for Service: ARTHUR S. CAVE & CO.
Patent & Trade Mark Attorneys
Level 10
10 Barrack Street
SYDNEY N.S.W. 2000
AUSTRALIA

**Complete Specification for the invention entitled AMINO ACID
DERIVATIVES.**

**The following statement is a full description of this invention
including the best method of performing it known to me:-**

Merck Patent Gesellschaft
mit beschränkter Haftung
6100 Darmstadt

Amino acid derivatives

The invention relates to new amino acid derivatives of the formula I



5 wherein

X

is H , $R^1-O-C_mH_{2m}-CO-$, $R^1-C_mH_{2m}-O-CO-$,
 $R^1-C_mH_{2m}-CO-$, R^1-SO_2- or
 $(R^1-C_mH_{2m}-(T)_x-C_rH_{2r})-L(R^7-C_pH_{2p})-C_tH_{2t}-CO-$,

10 Z

is 0 to 4 amino acid radicals attached to one another by a peptide linkage and selected from the group consisting of Abu, Ada, Ala, BAla, Arg, Asn, Asp, Bia, Cal, Dab, Gln, Glu, Gly, His, N(im)-alkyl-His, Ile, Leu, tert.-Leu, Lys, Met, α Nal, BNal, Nbg, Nle, Orn, Phe, Pro, Ser, Thr, Tic, Trp, Tyr and Val,

15


E

is 0 to 2 amino acid radicals attached to one another by a peptide linkage and selected from the group consisting of Abu, Ala, Cal, His, Ile, Leu, Met, Nle, Phe, Trp, Tyr and Val,

20 D

is $-CH_2-CHOH-CH_2OH$, $-C_2H_2-SO_2-R^{14}$, a phenyl- C_6H_5- , furyl- C_4H_3- , thienyl- C_4H_3- or pyridyl- C_5H_4- radical which is substituted by one or two $R^{14}-SO_2-$ groups or an $R^{14}-CO-$ group or an $(R^{14})_2-PO-$ group and, if appropriate, additionally by an Hal atom, or D is

25



- R^1 , R^3 , R^7 and R^8 are each H, A, Ar, Ar-alkyl, Het, Het-alkyl or cycloalkyl having 3-7 C atoms, cycloalkylalkyl having 4-11 C atoms, bicycloalkyl or tricycloalkyl having in each case 7-14 C atoms or bicycloalkylalkyl or tricycloalkylalkyl having in each case 8-18 C atoms, each of which is unsubstituted or monosubstituted or polysubstituted by A, AO and/or Hal,
- R^2 , R^5 and R^6 are each H or A,
- R^4 is (H, OH), (H, NH_2) or =O,
- R^9 is H, NH_2 , NHA or NA_2 ,
- R^{10} , R^{11} , R^{12} and R^{13} are each H, Hal, OH, OA, NH_2 , SH, SA, SO_2NH_2 , CF_3 , CN, COOH or COOA,
- R^{14} is OH, OA, NH_2 , NHA, NA_2 , NHCycloalkyl having 3-7 C atoms, N(cycloalkyl)₂ having 6-14 C atoms, pyrrolidino, piperidino, hexahydroazepino, morpholino, thiomorpholino, piperazino, N-A-piperazino, NHA₂ or NHHet,
- L is CH or N,
- T is O, S, NH or NA,
- n is 1 or 2,
- m, p, r and t are each 0, 1, 2, 3, 4 or 5,
- x is 0 or 1,
- y is 0, 1 or 2,
- z is 2, 3, 4, 5 or 6,
- Ar is phenyl which is unsubstituted or monosubstituted or polysubstituted by A, AO, Hal, CF_3 , OH, H_2NSO_2 and/or NH_2 , or unsubstituted naphthyl,
- Het is a saturated or unsaturated 5-membered or 6-membered heterocyclic radical which has 1-4 N, O and/or S atoms, which can be condensed with a benzene ring

and/or can be monosubstituted or polysubstituted by
A, AO, Hal, CF₃, HO, O₂N, carbonyl oxygen,
H₂N, HAN, A₂N, AcNH, AS, ASO, ASO₂, HOOC, AOO, C,
CN, H₂NCO, H₂NSO₂, ASO₂NH, Ar or Ar-alkenyl,
hydroxyalkyl and/or aminoalkyl having in each case
1-8 C atoms, and/or in which the N and/or S hetero-
atoms can also be oxidized,

Hal is F, Cl, Br or I,

Ac is A-CO-, Ar-CO- or A-NH-CO-,

-alkyl- is an alkylene group having 1-4 C atoms and

A is alkyl having 1-8 C atoms,

and wherein it is also possible for one or more -NA-CO-
groups to replace one or more -NH-CO- groups, and to salts
thereof.

Similar compounds are known from EP-A-77,028.

The invention was based on the object of finding
new compounds having valuable properties, especially com-
pounds which can be used for the preparation of medicaments.

It has been found that the compounds of the formula
I and their salts possess very valuable properties. Above
all, they inhibit the activity of human plasma renin. This
effect can be demonstrated, for example, by the method of
F. Fyhrquist et al., Clin. Chem. 22, 250-256 (1976). It is
remarkable that these compounds are very specific inhibitors
of renin; as a rule substantially higher concentrations of
these compounds are necessary for the inhibition of other
aspartylproteinases (for example pepsin and cathepsin D).

The compounds can be employed as active compounds
for medicaments in human and veterinary medicine, especially
for the prophylaxis and treatment of cardiac, circulatory
and vascular diseases, above all hypertension, cardiac in-
sufficiency and hyperaldosteronism. In addition, the com-
pounds can be used for diagnostic purposes in order to
determine, in the case of patients with hypertension or
hyperaldosteronism, the possible contribution made by the
renin activity towards maintaining the pathological state.

The abbreviations of amino acid radicals listed
above and below represent the radicals -NR'-R''-CO-, as a

rule -NH-CHR-CO-, (wherein R, R' and R" have the specific meaning known for each amino acid) of the following amino acids:

	Abu	2-aminobutyric acid
5	Ada	3-adamantylalanine
	Ala	alanine
	BAla	β -alanine
	Arg	arginine
	Asn	asparagine
10	Asp	aspartic acid
	Bia	3-(2-benzimidazolyl)-alanine
	Cal	3-cyclohexylalanine
	Dab	2,4-diaminobutyric acid
	Gln	glutamine
15	Glu	glutamic acid
	Gly	glycine
	His	histidine
	N(im)-alkyl-His	histidine which is substituted by A in the 1-position or 3-position of the imidazole ring
20	Ile	isoleucine
	Leu	leucine
	tert.-Leu	tert.-leucine
	Lys	lysine
25	Met	methionine
	α Nal	α -naphthylalanine
	β Nal	β -naphthylalanine
	Nbg	(2-norbornyl)-glycine
	Nle	norleucine
30	N-Me-His	N-methylhistidine
	N-Me-Phe	N-methylphenylalanine
	Orn	ornithine
	Phe	phenylalanine
	Pro	proline
35	Ser	serine
	Thr	threonine
	Tic	tetrahydroisoquinoline-1-carboxylic acid
	Trp	tryptophane
	Tyr	tyrosine

Val valine.

The abbreviations below also have the following meanings:

BOC tert.-butoxycarbonyl

5 imi-BOM benzyloxymethyl in the 1-position of the imidazole ring

CBZ benzyloxycarbonyl

DNP 2,4-dinitrophenyl

10 imi-DNP 2,4-dinitrophenyl in the 1-position of the imidazole ring

ETNC N-ethylcarbamoyl

ETOC ethoxycarbonyl

FMOC 9-fluorenylmethoxycarbonyl

IPNC N-isopropylcarbamoyl

15 IPOC isopropoxycarbonyl

MC morpholinocarbonyl

OMe methyl ester

OEt ethyl ester

PBB 4-phenyl-2-benzylbutyryl

20 POA phenoxyacetyl

DCCI dicyclohexylcarbodiimide

HOBt 1-hydroxybenzotriazole.

Insofar as the amino acids mentioned above can exist in several enantiomeric forms, all these forms and also
25 mixtures thereof (for example the DL-forms) are included in the above and following text, for example as a constituent of the compounds of the formula I. The L-forms are preferred. Where individual compounds are listed in the following text, the abbreviations of these amino acids relate in
30 each case to the L-form, unless anything to the contrary is expressly indicated.

The invention also relates to a process for the preparation of an amino acid derivative of the formula I and salts thereof, characterized in that it is liberated
35 from one of its functional derivatives by treatment with a solvolysing or hydrogenolysing agent, or in that a carboxylic acid of the formula II

wherein G^1 is (a) Z^1 ,
 (b) Z ,
 (c) $Z-W$,
 (d) $Z-W-E^1$,
 (e) $Z-W-E$ and

W is $-NR^2-CHR^3-CR^4-(CHR^5)_n-CO-$

is reacted with an amino compound of the formula III
 $H-G^2$ III

wherein G^2 is (a) $-Z^2-W-E-NR^6-D$,
 (b) $-W-E-NR^6-D$,
 (c) $-E-NR^6-D$,
 (d) $-E^2-NR^6-D$,
 (e) $-NR^6-D$,

E^1 and E^2 are each one amino acid radical selected from the group consisting of Abu, Ala, Cal, His, Ile, Leu, Met, Nle, Phe, Trp, Tyr and Val in such a manner that $E^1 + E^2$ together are E,

Z^1 and Z^2 are each 1 to 3 amino acid radicals selected from the group consisting of Abu, Ada, Ala, BAla, Arg, Asn, Asp, Bia, Cal, Dab, Gln, Glu, Gly, His N(im)-alkyl-His, Ile, Leu, tert -Leu, Lys, Met, α Nal, BNal, Nbg, Nle, Orn, Phe, Pro, Ser, Thr, Tic, Trp, Tyr and Val in such a manner that $Z^1 + Z^2$ together are Z, and in that, if appropriate, a functionally modified amino and/or hydroxyl group in a compound of the formula I is liberated by treatment with solvolysing or hydrogenolysing agents and/or, in order to prepare a compound of the Formula I wherein R^4 is (H, OH) or (H, NH_2), an aminoketo acid derivative of the formula I wherein $R^4 = 0$ is reduced or reductively aminated and/or a radical D is converted into another radical D by treatment with esterifying, solvolysing or reducing agents and/or a compound of the formula I is converted into one of its salts by treatment with an acid.

In the preceding and following text, the radicals or parameters X, Z, E, D, R^1 to R^{14} , L, T, m, n, p, r, t, x, y, z, Ar, Het, Hal, Ac, A, G^1 , G^2 , E^1 , E^2 , Z^1 , Z^2 and W have the meanings indicated in the formulae I, II or III, unless anything to the contrary is expressly indicated.

In the formulae above, A has 1 - 8, preferably 1, 2, 3 or 4, C atoms. A is preferably methyl and also ethyl, propyl, isopropyl, butyl, isobutyl, sec.-butyl or tert.-butyl, and also pentyl, 1-, 2- or 3-methylbutyl, 1,1-, 1,2- or 2,2-dimethylpropyl, 1-ethylpropyl, hexyl, 1-, 2-, 3- or

4-methylpentyl, 1,1-, 1,2-, 1,3-, 2,2-, 2,3- or 3,3-dimethylbutyl, 1-ethylbutyl, 2-ethylbutyl, 1-ethyl-1-methylpropyl, 1-ethyl-2-methylpropyl, 1,1,2-trimethylpropyl, 1,2,2-trimethylpropyl, heptyl or octyl.

5 Cycloalkyl is preferably cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl or cycloheptyl, but is also, for example, 1-, 2- or 3-methylcyclopentyl or 1-, 2-, 3- or 4-methylcyclohexyl.

10 Accordingly, cycloalkylalkyl is preferably cyclopropylmethyl, 2-cyclopropylethyl, cyclobutylmethyl, 2-cyclobutylethyl, cyclopentylmethyl, 2-cyclopentylethyl, cyclohexylmethyl or 2-cyclohexylethyl, but is also, for example, 1-, 2- or 3-methylcyclopentylmethyl or 1-, 2-, 3- or 4-methylcyclohexylmethyl.

15 Bicycloalkyl is preferably 1-decalyl, 2-decalyl, 2-bicyclo[2,2,1]heptyl or 6,6-dimethyl-2-bicyclo[3,1,1]heptyl.

Tricycloalkyl is preferably 1-adamantyl.

Hal is preferably F, Cl or Br, but also I.

20 Ac is preferably A-CO-, such as acetyl, propionyl or butyryl, Ar-CO-, such as benzoyl, o-, m- or p-methoxybenzoyl or 3,4-dimethoxybenzoyl, or A-NH-CO-, such as N-methylcarbamoyl or N-ethylcarbamoyl.

25 Ar is preferably phenyl and also preferably o-, m- or p-tolyl, o-, m- or p-ethylphenyl, o-, m- or p-methoxyphenyl, o-, m- or p-fluorophenyl, o-, m- or p-chlorophenyl, o-, m- or p-bromophenyl, o-, m- or p-iodophenyl, o-, m- or p-trifluoromethylphenyl, o-, m- or p-hydroxyphenyl, o-, m- or p-sulfamoylphenyl, 2,3-, 2,4-, 2,5-, 2,6-, 3,4- or 3,5-dimethoxyphenyl, 3,4,5-trimethoxyphenyl, o-, m- or p-amino-
30 phenyl, 1-naphthyl or 2-naphthyl.

 Accordingly, Ar-alkyl is preferably benzyl, 1-phenylethyl, 2-phenylethyl, o-, m- or p-methylbenzyl, 1-o-, -m- or -p-tolylolethyl, 2-o-, -m- or -p-tolylolethyl, o-, m- or p-ethylbenzyl, 1-o-, -m- or -p-ethylphenylethyl, 2-o-, -m- or -p-ethylphenylethyl, o-, m- or p-methoxybenzyl, 1-o-, -m- or -p-methoxyphenylethyl, 2-o-, -m- or -p-methoxyphenyl-
35 ethyl, o-, m- or p-fluorobenzyl, 1-o-, -m- or p-fluoro-

phenylethyl, 2-o-, -m- or -p-fluorophenylethyl, o-, m- or
p-chlorobenzyl, 1-o-, -m- or -p-chlorophenylethyl, 2-o-,
-m- or -p-chlorophenylethyl, o-, m- or p-bromobenzyl, 1-o-,
-m- or -p-bromophenylethyl, 2-o-, -m- or -p-bromophenyl-
5 ethyl, o-, m- or p-iodobenzyl, 1-o-, -m- or -p-iodophenyl-
ethyl, 2-o-, -m- or -p-iodophenylethyl, o-, m- or p-tri-
fluoromethylbenzyl, o-, m- or p-hydroxybenzyl, 2,3-, 2,4-,
2,5-, 2,6-, 3,4- or 3,5-dimethoxybenzyl, 3,4,5-trimethoxy-
benzyl, o-, m- or p-aminobenzyl, 1-naphthylmethyl or 2-naph-
10 thylmethyl.

Het is preferably 2-furyl, 3-furyl, 2-thienyl, 3-
thienyl, 1-, 2- or 3-pyrryl, 1-, 2-, 4- or 5-imidazolyl, 1-,
3-, 4- or 5-pyrazolyl, 2-, 4- or 5-oxazolyl, 3-, 4- or 5-
isoxazolyl, 2-, 4- or 5-thiazolyl, 3-, 4- or 5-isothiazolyl,
15 2-, 3- or 4-pyridyl or 2-, 4-, 5- or 6-pyrimidyl, and is
also preferably 1,2,3-triazol-1-yl, -4-yl or -5-yl, 1,2,4-
triazol-1-yl, -3-yl or -5-yl, 1-tetrazolyl, 5-tetrazolyl,
1,2,3-oxadiazol-4-yl, 1,2,3-oxadiazol-5-yl, 1,2,4-oxadiazol-
3-yl, 1,2,4-oxadiazol-5-yl, 1,3,4-thiadiazol-2-yl, 1,3,4-
20 thiadiazol-5-yl, 1,2,4-thiadiazol-3-yl, 1,2,4-thiadiazol-5-
yl, 2,1,5-thiadiazol-3-yl, 2,1,5-thiadiazol-4-yl, 2-, 3-,
4-, 5- or 6-2H-thiopyranyl, 2-, 3- or 4-4H-thiopyranyl, 3-
pyridazinyl, 4-pyridazinyl, pyrazinyl, 2-, 3-, 4-, 5-, 6-
or 7-benzofuryl, 2-, 3-, 4-, 5-, 6- or 7-benzothienyl, 1-,
25 2-, 3-, 4-, 5-, 6- or 7-indolyl, 1-, 2-, 3-, 4-, 5-, 6- or
7-isoindolyl, 1-, 2-, 4- or 5-benzimidazolyl, 1-, 3-, 4-,
5-, 6- or 7-benzopyrazolyl, 2-, 4-, 5-, 6- or 7-benzoxazolyl,
3-, 4-, 5-, 6- or 7-benzisoxazolyl, 2-, 4-, 5-, 6- or 7-
benzthiazolyl, 2-, 4-, 5-, 6- or 7-benzisothiazolyl, 4-,
30 5-, 6- or 7-benz-2,1,3-oxadiazolyl, 2-, 3-, 4-, 5-, 6-, 7-
or 8-quinolyl, 1-, 3-, 4-, 5-, 6-, 7- or 8-isoquinolyl, 1-,
2-, 3-, 4- or 9-carbazolyl, 1-, 2-, 3-, 4-, 5-, 6-, 7-, 8-
or 9-acridinyl, 3-, 4-, 5-, 6-, 7- or 8-cinnolyl, 2-, 4-,
5-, 6-, 7- or 8-quinazolyl. The heterocyclic radicals can
35 also be partly or completely hydrogenated. Het can, there-
fore, also be, for example: 2,3-dihydro-2-, -3-, -4- or -5-
furyl, 2,5-dihydro-2-, -3-, -4- or 5-furyl, tetrahydro-2-
furyl, tetrahydro-3-furyl, tetrahydro-2-thienyl, tetrahydro-

3-thienyl, 2,3-dihydro-1-, -2-, -3-, -4- or -5-pyrryl, 2,5-dihydro-1-, -2-, -3-, -4- or -5-pyrryl, 1-, 2- or 3-pyrrolidinyl, tetrahydro-1-, -2- or -4-imidazolyl, 2,3-dihydro-1-, -2-, -3-, -4- or -5-pyrazolyl, 2,5-dihydro-1-, -2-, -3-, -4- or 5-pyrazolyl, tetrahydro-1-, -3- or -4-pyrazolyl, 1,4-dihydro-1-, -2-, -3- or -4-pyridyl, 1,2,3,4-tetrahydro-1-, -2-, -3-, -4-, -5- or -6-pyridyl, 1,2,3,6-tetrahydro-1-, -2-, -3-, -4-, -5- or -6-pyridyl, 1-, 2-, 3- or 4-piperidinyl, 2-, 3- or 4-morpholinyl, tetrahydro-2-, -3- or -4-pyranyl, 1,4-dioxanyl, 1,3-dioxan-2-, -4- or -5-yl, hexahydro-1-, -3- or -4-pyridazinyl, hexahydro-1-, -2-, -4- or -5-pyrimidyl, 1-, 2- or 3-piperazinyl, 1,2,3,4-tetrahydro-1-, -2-, -3-, -4-, -5-, -6-, -7- or -8-quinolyl, 1,2,3,4-tetrahydro-1-, -2-, -3-, -4-, -5-, -6-, -7- or 8-isoquinolyl.

The heterocyclic radicals can also be substituted as indicated. Het can, therefore, preferably also be: 2-amino-4-thiazolyl, 4-carboxy-2-thiazolyl, 4-carbamoyl-2-thiazolyl, 4-(2-aminoethyl)-2-thiazolyl, 2-amino-5,6-dimethyl-3-pyrazinyl or 4-carbamoylpiperidino, and also, for example, 3-, 4- or 5-methyl-2-furyl, 2-, 4- or 5-methyl-3-furyl, 2,4-dimethyl-3-furyl, 5-nitro-2-furyl, 5-styryl-2-furyl, 3-, 4- or 5-methyl-2-thienyl, 2-, 4- or 5-methyl-3-thienyl, 3-methyl-5-tert.-butyl-2-thienyl, 5-chloro-2-thienyl, 5-phenyl-2- or -3-thienyl, 1-, 3-, 4- or 5-methyl-2-pyrryl, 1-methyl-4-nitro-2-pyrryl, 1-methyl-5-nitro-2-pyrryl, 3,5-dimethyl-4-ethyl-2-pyrryl, 4-methyl-5-pyrazolyl, 5-methyl-3-isoxazolyl, 3,4-dimethyl-5-isoxazolyl, 4-methyl-2-thiazolyl, 5-methyl-2-thiazolyl, 2-methyl-4-thiazolyl, 5-methyl-4-thiazolyl, 2-methyl-5-thiazolyl, 4-methyl-5-thiazolyl, 2,4-dimethyl-5-thiazolyl, 3-, 4-, 5- or 6-methyl-2-pyridyl, 2-, 4-, 5- or 6-methyl-3-pyridyl, 2- or 3-methyl-4-pyridyl, 3-, 4-, 5- or 6-chloro-2-pyridyl, 2-, 4-, 5- or 6-chloro-3-pyridyl, 2-chloro-4-pyridyl, 3-chloro-4-pyridyl, 2,6-dichloropyridyl, 2-hydroxy-3-, -4-, -5- or -6-pyridyl (= 1H-2-pyridon-3-, -4-, -5- or -6-yl), 5-phenyl-1H-2-pyridon-3-yl, 5-p-methoxyphenyl-1H-2-pyridon-3-yl, 2-methyl-3-hydroxy-4-hydroxymethyl-5-pyridyl, 2-hydroxy-4-amino-6-methyl-3-pyridyl, 3-N'-methylureido-1H-4-pyridon-5-yl, 4-

methyl-2-pyrimidyl, 4,6-dimethyl-2-pyrimidyl, 2-, 5- or 6-methyl-4-pyrimidyl, 2,6-dimethyl-4-pyrimidyl, 2,6-dihydroxy-4-pyrimidyl, 5-chloro-2-methyl-4-pyrimidyl, 2-methyl-4-amino-5-pyrimidyl, 3-methyl-2-benzofuryl, 2-ethyl-3-benzofuryl, 7-methyl-2-benzothienyl, 1-, 2-, 4-, 5-, 6- or 7-methyl-3-indolyl, 1-methyl-5- or -6-benzimidazolyl, 1-ethyl-5-benzimidazolyl, 1-ethyl-6-benzimidazolyl and 3-, 4-, 5-, 6-, 7- or 8-hydroxy-2-quinolyl.

5 R^1 and R^7 are preferably A, especially methyl, ethyl, propyl, isopropyl, butyl, isobutyl or tert.-butyl, and also, preferably, cyclopropyl, cyclopentyl, cyclohexyl, phenyl, benzyl, pyrrolidino, piperidino or morpholino.

R^2 , R^5 and R^6 are preferably H or methyl, and also ethyl, propyl, isopropyl, butyl or isobutyl.

15 R^3 is preferably cyclohexylmethyl, and also preferably A, especially methyl, ethyl, propyl, isopropyl, butyl, isobutyl, sec.-butyl, pentyl, isopentyl (3-methylbutyl) or 2-methylbutyl, phenyl, benzyl, p-chlorobenzyl, 2-cyclohexylethyl, bicyclo[2,2,1]heptyl-2-methyl or 6,6-dimethylbicyclo[3,1,1]heptyl-2-methyl.

20 R^4 is preferably (H, OH).

R^8 is preferably isopropyl, isobutyl, sec.-butyl or benzyl, and also preferably H, methyl, ethyl, propyl, butyl or cyclohexylmethyl.

25 R^9 is preferably H or NH_2 .

R^{10} and R^{13} are preferably H.

R^{11} is preferably H or SO_2NH_2 .

R^{12} is preferably H or Cl.

30 R^{14} is preferably NH_2 , $NHCH_3$ or $N(CH_3)_2$, and also preferably 2-thiazolylamino, 3-isoxazolylamino, 5-methyl-3-isoxazolylamino, 3,4-dimethyl-5-isoxazolylamino, 2-pyrimidylamino, 4-methyl-2-pyrimidylamino, 4,6-dimethyl-2-pyrimidylamino or 2,6-dimethyl-4-pyrimidylamino.

L is preferably CH.

35 T is preferably O or S.

The parameters m, p, r and t are preferably 0, 1 or 2; n is preferably 1; x is preferably 0; y is preferably 0 or 1; and z is preferably 2.

X is preferably H, POA, alkoxycarbonyl, such as ETOC, IPOC or BOC, CBZ, alkanoyl, such as acetyl, propionyl, butyryl or isobutyryl, cycloalkylcarbonyl, such as cyclopentylcarbonyl or cyclohexylcarbonyl, aroyl, such as benzoyl, arylalkanoyl, such as phenylacetyl, 2-phenylpropionyl, 3-phenylpropionyl, 4-phenylbutyryl, 2-benzyl-3-phenylpropionyl, PBB, 2-(2-phenylethyl)-4-phenylbutyryl, 2-(2-naphthylmethyl)-4-phenylbutyryl, 2-o-, -m- or -p-fluorophenylpropionyl, 3-o-, -m- or -p-fluorophenylpropionyl, 2-o-, -m- or -p-chlorophenylpropionyl, or 3-o-, -m- or -p-chlorophenylpropionyl, or cycloalkylalkanoyl, such as cyclohexylacetyl, 2-cyclohexylpropionyl or 3-cyclohexylpropionyl, or N-alkylcarbamoyl, such as ETNC or IPNC or MC. Radicals X which are particularly preferred are BOC and MC, and also ETOC, IPOC, ETNC, IPNC and PBB, and also H, POA, 4-phenylbutyryl, 2-benzyl-3-phenylpropionyl, 2-(2-phenylethyl)-4-phenylbutyryl, 2-(2-naphthylmethyl)-4-phenylbutyryl and CBZ. A further group of particularly preferred radicals X corresponds to the formula $R^0-CH(CH_2C_6H_5)-CO-$ wherein R^0 is pyrrolidino, piperidino, morpholino, alkyl, alkoxy or alkylthio each of which has 1-8 C atoms.

Z is preferably 2, but also 0 or 1 and also 3 or 4, amino acid radicals which are attached to one another by a peptide linkage, in particular one of the groups Gly, His, Phe-Gly, Phe-His, Pro-Phe-His or His-Pro-Phe-His, and also preferably one of the groups Abu, Ada, Asn, Bia, Cal, Gln, N-(im)-methyl-His, Leu, α Nal, BNal, Nle, Phe, Trp, Tyr, Abu-His, Ada-His, Ala-His, Ala-Phe, Arg-His, Asn-His, Bia-His, Cal-His, Dab-His, Glu-His, Gly-His, His-His, Ile-His, Leu-His, tert.-Leu-His, Lys-His, Met-His, α Nal-His, BNal-His, Nbg-His, Nle-His, (N-Me-His)-His, (N-Me-Phe)-His, Orn-His, Phe-Abu, Phe-Ada, Phe-Ala, Phe-Arg, Phe-Asn, Phe-Bia, Phe-Cal, Phe-Dab, Phe-Gln, Phe-Glu, Phe-(N-im-methyl-His), Phe-Ile, Phe-Leu, Phe-tert.Leu, Phe-Lys, Phe-Met, Phe- α -Nal, Phe-BNal, Phe-Nbg, Phe-Nle, Phe-(N-Me-His), Phe-(N-Me-Phe), Phe-Orn, Phe-Phe, Phe-Pro, Phe-Ser, Phe-Thr, Phe-Tic, Phe-Trp, Phe-Tyr, Phe-Val, Pro-His, Ser-His, Thr-His, Tic-His, Trp-His, Tyr-His, Val-His, and also Ada-Phe-His, Pro-Ala-

His, Pro-Ala-Phe, Pro-Phe-Ala, Pro-Phe-Phe, His-Pro-Ala-His,
and also Pro-Abu-His, Pro-Ada-His, Pro-Arg-His, Pro-Asn-His,
Pro-Bia-His, Pro-Dab-His, Pro-Glu-His, Pro-His-His, Pro-
Ile-His, Pro-Leu-His, Pro-tert.-Leu-His, Pro-Lys-His, Pro-
5 Met-His, Pro-Nbg-His, Pro-Nle-His, Pro-(N-Me-His)-His,
Pro-(N-Me-Phe)-His, Pro-Orn-His, Pro-Phe-Abu, Pro-Phe-Ada,
Pro-Phe-Arg, Pro-Phe-Asn, Pro-Phe-Bia, Pro-Phe-Dab, Pro-
Phe-Gln, Pro-Phe-Glu, Pro-Phe-(N-im-methyl-His), Pro-Phe-
Ile, Pro-Phe-Leu, Pro-Phe-tert.-Leu, Pro-Phe-Lys, Pro-Phe-
10 Met, Pro-Phe-Nbg, Pro-Phe-Nle, Pro-Phe-(N-Me-His), Pro-Phe-
(N-Me-Phe), Pro-Phe-Orn, Pro-Phe-Pro, Pro-Phe-Ser, Pro-Phe-
Thr, Pro-Phe-Tic, Pro-Phe-Trp, Pro-Phe-Tyr, Pro-Phe-Val,
Pro-Pro-His, Pro-Ser-His, Pro-Thr-His, Pro-Tic-His, Pro-
Trp-His, Pro-Tyr-His, Pro-Val-His, His-Pro-Abu-His, His-
15 Pro-Ada-His, His-Pro-Arg-His, His-Pro-Asn-His, His-Pro-Bia-
His, His-Pro-Dab-His, His-Pro-Glu-His, His-Pro-His-His,
His-Pro-Ile-His, His-Pro-Leu-His, His-Pro-tert.-Leu-His,
His-Pro-Lys-His, His-Pro-Met-His, His-Pro-Nbg-His, His-Pro-
Nle-His, His-Pro-(N-Me-His)-His, His-Pro-(N-Me-Phe)-His,
20 His-Pro-Orn-His, His-Pro-Phe-Abu, His-Pro-Phe-Ada, His-Pro-
Phe-Arg, His-Pro-Phe-Asn, His-Pro-Phe-Bia, His-Pro-Phe-Dab,
His-Pro-Phe-Gln, His-Pro-Phe-Glu, His-Pro-Phe(N-im-methyl-
His), His-Pro-Phe-Ile, His-Pro-Phe-Leu, His-Pro-Phe-tert.-
Leu, His-Pro-Phe-Lys, His-Pro-Phe-Met, His-Pro-Phe-Nbg,
25 His-Pro-Phe-Nle, His-Pro-Phe-(N-Me-His), His-Pro-Phe-(N-Me-
Phe), His-Pro-Phe-Orn, His-Pro-Phe-Pro, His-Pro-Phe-Ser,
His-Pro-Phe-Thr, His-Pro-Phe-Tic, His-Pro-Phe-Trp, His-Pro-
Phe-Tyr, His-Pro-Phe-Val, His-Pro-Pro-His, His-Pro-Ser-His,
His-Pro-Thr-His, His-Pro-Tic-His, His-Pro-Trp-His, His-Pro-
30 Tyr-His, His-Pro-Val-His.

If X is one of the groups $R^0-CH(CH_2C_6H_5)-CO-$,
Z is preferably Gly or His.

E is preferably absent or is preferably Ile or Leu,
and also preferably Abu, Cal, Met or Nle.

35 D is preferably $-CH_2-CHOH-CH_2OH$, $-(CH_2)_2SO_2NH_2$, $-o-$,
 $-m-$ or especially $-p-C_6H_4-SO_2NH_2$, $-o-$, $-m-$ or especially $-p-$
 $CH_2C_6H_4-SO_2NH_2$, $-o-$, $-m-$ or especially $-p-C_6H_4-SO_2NHCH_3$, $-o-$,
 $-m-$ or especially $-p-C_6H_4-SO_2N(CH_3)_2$, $-o-$, $-m-$ or especially

-p-C₆H₄SO₂NHHet, -o-, -m- or especially p-C₆H₄CONH₂, 5-sulfamoyl-2-pyridyl, 2-sulfamoyl-5-thienyl or 3-sulfamoyl-5-thienyl, 3H-quinazolin-4-on-2-yl-CHR⁸, 3-amino-3H-quinazolin-4-on-2-yl-CHR⁸, 6-aminosulfonyl-7-chloro-3H-quinazolin-4-on-2-yl-CHR⁸ or 3-amino-6-aminosulfonyl-7-chloro-3H-quinazolin-4-on-2-yl-CHR⁸.

The group W is preferably -NH-CHR³-CHOH-CH₂-CO-, especially -NH-CH(cyclohexylmethyl)-CHOH-CH₂-CO- ("AHCP", derived from 4-amino-3-hydroxy-5-cyclohexylpentanoic acid) and also -NH-CH(CH₂CH₂-cyclohexyl)-CHOH-CH₂-CO- ("AHCH"; derived from 4-amino-3-hydroxy-6-cyclohexylhexanoic acid), -NH-CH(isobutyl)-CHOH-CH₂-CO- ("Sta"; derived from statin) or -NH-CH(benzyl)-CHOH-CH₂-CO- ("AHPP"; derived from 4-amino-3-hydroxy-5-phenylpentanoic acid). The group W is also preferably -NH-CHR³-CH(NH₂)-CH₂-CO-, especially -NH-CH(cyclohexylmethyl)-CH(NH₂)-CH₂-CO- ("DACP"; derived from 3,4-diamino-5-cyclohexylpentanoic acid), -NH-CH(CH₂CH₂-cyclohexyl)-CH(NH₂)-CH₂-CO- ("DACH"; derived from 3,4-diamino-6-cyclohexylhexanoic acid), -NH-CH(isobutyl)-CH(NH₂)-CH₂-CO- ("DAMH"; derived from 3,4-diamino-6-methylheptanoic acid) or -NH-CH(benzyl)-CH(NH₂)-CH₂-CO- ("DAPP"; derived from 3,4-diamino-5-phenylpentanoic acid).

The group W has at least one chiral centre. Further chiral centres can be present in the groups X, R⁶ and D. The compounds of the formula I can, therefore, exist in various forms - optically inactive or optically active. The formula I embraces all these forms. If W is -NH-CHR³-CR⁴-CH₂-CO- in which R⁴ is (H, OH) or (H, NH₂), the 3S-hydroxy-4S-amino enantiomers or 3S,4S-diamino enantiomers are preferred. Unless anything to the contrary is indicated in the designation of individual substances, the abbreviations AHCP, AHCH, Sta, AHPP, DACP, DACH, DAMH and DAPP relate in each case to the 3S,4S-forms.

Accordingly, the invention relates particularly to compounds of the formula I in which at least one of the radicals mentioned has one of the preferred meanings indicated above. Some preferred groups of compounds can be

expressed by means of the partial formulae Ia to Il following, which correspond to the formula I, but in which

in Ia X is H, BOC or $R^0\text{-CH(CH}_2\text{C}_6\text{H}_5\text{)-CO-}$;

in Ib X is BOC;

5 in Ic Z is Gly, His, Phe-Gly or Phe-His;

in Id Z is Phe-Gly or Phe-His;

in Ie Z is Phe-His;

in If $\text{-NR}^2\text{-CHR}^3\text{-CR}^4\text{-(CHR}^5\text{)}_n\text{-CO-}$ (= W) is AHCP;

in Ig E is absent;

in Ih R^6 is H;

10 in Ii X is H, BOC or $R^0\text{-CH(CH}_2\text{C}_6\text{H}_5\text{)-CO-}$,

Z is Gly, His, Phe-Gly or Phe-His,

W is AHCP,

R^6 is H and

E is absent;

15 in Ij X is BOC,

Z is Phe-His or Phe-Gly,

W is AHCP,

R^6 is H and

E is absent;

20 in Ik X is BOC and

Z is Phe-Gly or Phe-His; and

in Il X is $R^0\text{-CH(CH}_2\text{C}_6\text{H}_5\text{)-CO-}$ and

Z is Gly or His.

Compounds of the following partial formulae are particularly preferred:

I* and Ia* to Il*, which correspond to the formulae I and Ia to Il, but in which

D is $\text{-CH}_2\text{-CHOH-CH}_2\text{OH}$, $\text{-p-C}_6\text{H}_4\text{-SO}_2\text{NH}_2$ or

30 $3\text{-R}^9\text{-5-R}^{10}\text{-6-R}^{11}\text{-7-R}^{12}\text{-8-R}^{13}\text{-3H-quinazolin-4-on-2-yl-CHR}^8\text{-}$;

I' and Ia' to Il', which correspond to the formulae I and Ia to Il, but in which

D is $\text{-CH}_2\text{-CHOH-CH}_2\text{OH}$;

35 I'' and Ia'' to Il'', which correspond to the formulae I and Ia to Il, but in which

D is $\text{-p-C}_6\text{H}_4\text{-SO}_2\text{NH}_2$;

I''' and Ia''' to Il''', which correspond to the formulae I and Ia to Il, but in which

on-2-yl-CHR⁸-,

R⁸ is H, alkyl having 1-4 C atoms or benzyl,

R⁹ is H or NH₂,

R¹⁰ and R¹³ are H,

5 R¹¹ is H or SO₂NH₂ and

R¹² is H or Cl;

I⁰ and Ia⁰ to Il⁰, which correspond to the formulae I and Ia to Il, but in which

D is 3H-quinazolin-4-on-2-yl-CHR⁸- or

10 3-amino-3H-quinazolin-4-on-2-yl-CHR⁸- and

R⁸ is H, alkyl having 1-4 C atoms or benzyl; and I⁰⁰ and Ia⁰⁰ to Il⁰⁰, which correspond to the formulae I and Ia to Il, but in which

D is 3H-quinazolin-4-on-1-yl-CHR⁸- or

15 3-amino-3H-quinazolin-4-on-2-yl-CHR⁸- and

R⁸ is sec.-butyl or isobutyl.

The compounds of the formula I and also the starting materials for their preparation are, incidentally, prepared by methods known per se, such as are described in the literature (for example in the standard works, such as Houben-Weyl, Methoden der organischen Chemie ("Methods of Organic Chemistry"), Georg-Thieme-Verlag, Stuttgart; and also EP-A-45,665, EP-A-77,028, EP-A-77,029 and EP-A-81,783), in particular under reaction conditions which are known and suitable for the reactions mentioned. In this regard it is also possible to make use of variants which are known per se, but are not mentioned here in detail.

If desired, the starting materials can also be formed in situ, so that they are not isolated from the reaction mixture, but are immediately reacted further to give the compounds of the formula I.

The compounds of the formula I are preferably obtained by liberating them from their functional derivatives by solvolysis, in particular hydrolysis, or by hydrogenolysis.

Preferred starting materials for the solvolysis or hydrogenolysis are those which otherwise correspond to the formula I, but, instead of one or more free amino and/or

hydroxyl groups, contain corresponding protected amino and/or hydroxyl groups, preferably groups of this type which, instead of an H atom attached to an N atom, carry an amino protective group, for example those which correspond to the formula I, but, instead of an His group, contain an N-(im)-
 5 R^{15} -His group (wherein R^{15} is an amino protective group, for example BOM or DNP), or those of the formula $X-Z-NR^2-CHR^3-CH(NHR^{15})-(CHR^5)_n-CO-E-NR^6-D$.

Preferred starting materials are also those which,
 10 instead of the H atom of a hydroxyl group, carry a hydroxyl protective group, for example those of the formula $X-Z-NR^2-CHR^3-CHOR^{16}-(CHR^5)_n-CO-E-NR^6-D$, wherein R^{16} is a hydroxyl protective group.

It is also possible for several - identical or
 15 different - protected amino and/or hydroxyl groups to be present in the molecule of the starting material. If the protective groups present are different from one another, they can in many cases be split off selectively.

The term "amino protective group" is generally
 20 known and relates to groups which are suitable for protecting (blocking) an amino group against chemical reactions, but which can be removed readily after the desired chemical reaction has been carried out at another point on the molecule. Typical representatives of groups of this kind are,
 25 in particular, unsubstituted or substituted acyl, aryl (for example DNP), aralkoxymethyl (for example BOM) or aralkyl (for example benzyl, 4-nitrobenzyl or triphenylmethyl) groups. Since the amino protective groups are removed after the desired reaction (or reaction sequence), their nature
 30 and size is, incidentally, not critical; groups having 1-20, in particular 1-8, C atoms are, however, preferred. In the context of the present process, the term "acyl group", is to be understood in the widest sense. It embraces acyl groups derived from aliphatic, araliphatic, aromatic or
 35 heterocyclic carboxylic acids or sulfonic acids and also, in particular, alkoxycarbonyl, aryloxycarbonyl and, above all, aralkoxycarbonyl groups. Examples of acyl groups of this type are alkanoyl, such as acetyl, propionyl or buty-

ryl; aralkanoyl, such as phenylacetyl; aroyl, such as benzoyl or toluyyl; aryloxyalkanoyl, such as POA; alkoxy-carbonyl, such as methoxycarbonyl, ETOC, 2,2,2-trichloro-ethoxycarbonyl, BOC or 2-iodoethoxycarbonyl; and aralkyl-oxycarbonyl, such as CBZ ("carbobenzoxyl"), 4-methoxybenzyl-oxycarbonyl or FMOC. Preferred amino protective groups are DNP and BOM, and also CBZ, FMOC, benzyl and acetyl.

The term "hydroxyl protective group" is also generally known and relates to groups which are suitable for protecting a hydroxyl group against chemical reactions, but which can be removed readily after the desired chemical reaction has been carried out at another point in the molecule. Typical representatives of such groups are the unsubstituted or substituted aryl, aralkyl or acyl groups mentioned above, and also alkyl groups. The nature and size of the hydroxyl protective groups is not critical, since they are removed again after the desired chemical reaction or reaction sequence; groups having 1-20, in particular 1-10, C atoms are preferred. Examples of hydroxyl protective groups are, inter alia: benzyl, p-nitrobenzoyl, p-toluenesulfonyl and acetyl, benzyl and acetyl being particularly preferred.

The functional derivatives of the compounds of the formula I to be used as starting materials can be prepared by customary methods of amino acid and peptide synthesis, such as are described, for example, in the standard works and patent applications mentioned, and also, for example, by the Merrifield solid phase method.

The liberation of the compounds of the formula I from their functional derivatives is effected - depending on the protective group used - with, for example, strong acids, preferably trifluoroacetic acid or perchloric acid, but also other strong inorganic acids, such as hydrochloric acid or sulfuric acid, strong organic carboxylic acids, such as trichloroacetic acid, or sulfonic acids, such as benzenesulfonic or p-toluenesulfonic acid. The presence of an additional inert solvent is possible, but not always necessary. Suitable inert solvents are preferably organic

solvents, for example carboxylic acids, such as acetic acid, ethers, such as tetrahydrofuran or dioxane, amides, such as dimethylformamide (DMF), halogenated hydrocarbons, such as methylene chloride, and also alcohols, such as methanol, ethanol or isopropanol, and also water. Mixtures of the abovementioned solvents are also suitable. Trifluoroacetic acid is preferably used in excess without the addition of a further solvent; perchloric acid is used in the form of a mixture of acetic acid and 70% perchloric acid in a 9 : 1 ratio. The reaction temperatures for the cleavage are preferably between about 0 and about 50°; the reaction is preferably carried out between 15 and 30° (room temperature).

The BOC group can be split off, for example, preferably by means of 40% trifluoroacetic acid in methylene chloride or by means of about 3 N to 5 N HCl in dioxane at 15-30°, while the Fmoc group can be split off by means of an approximately 5 to 20% solution of dimethylamine, diethylamine or piperidine in DMF at 15-30°. Splitting off the DNP group is possible, for example, also by means of an approximately 3 to 10% solution of 2-mercaptoethanol in DMF/water at 15-30°.

Protective groups which can be removed by hydrogenolysis (for example BOM, CBZ or benzyl) can be split off, for example, by treatment with hydrogen in the presence of a catalyst (for example a noble metal catalyst such as palladium, preferably on a support such as charcoal). Suitable solvents for this reaction are those indicated above, in particular, for example, alcohols, such as methanol or ethanol, or amides, such as DMF. As a rule, the hydrogenolysis is carried out at temperatures between about 0 and 100° and pressures between about 1 and 200 bar, preferably at 20-30° and 1-10 bar. Hydrogenolysis of the CBZ group can be effected readily, for example, over 5 to 10% Pd/C in methanol at 20-30°.

Compounds of the formula I can also be obtained by direct peptide synthesis from a carboxylic acid component (formula II) and an amine component (formula III).

of suitable carboxylic acid components are those of the partial formulae $X-Z-OH$, $X-Z-W-OH$ or $X-Z-W-E-OH$, while suitable amine components are those of the partial formulae $H-W-E-NR^6-D$, $H-E-NR^6-D$ or $H-NR^6-D$. The peptide bond can, however, also be made within the group Z or E ; in this case a carboxylic acid of the formulae $X-Z^1-OH$ or $H-Z-W-E^1-OH$ with an amino compound of the formula $H-Z^2-W-E-NR^6-D$ or $H-E^2-NR^6-D$, respectively, $Z^1 + Z^2$ being Z or $E^1 + E^2$ being E , respectively. This reaction is preferably carried out by customary methods of peptide synthesis, such as are described, for example, in Houben-Weyl, loc. cit., volume 15/II, pages 1 to 806 (1974).

The reaction is preferably carried out in the presence of a dehydrating agent, for example a carbodiimide, such as DCCI or dimethylaminopropylethylcarbodiimide, and also propanephosphonic anhydride (compare Angew. Chem. 92, 129 (1980)), diphenylphosphoryl azide or 2-ethoxy-N-ethoxycarbonyl-1,2-dihydroquinoline, in an inert solvent, for example a halogenated hydrocarbon, such as methylene chloride, an ether, such as tetrahydrofuran or dioxane, an amide, such as DMF or dimethylacetamide, or a nitrile, such as acetonitrile, at temperatures between about -10 and 40° , preferably between 0 and 30° .

Instead of II or III, it is also possible to employ in the reaction suitable reactive derivatives of these compounds, for example compounds in which reactive groups are blocked in the meantime by protective groups. The amino acid derivatives III can, for example, be used in the form of their activated esters, which are preferably formed in situ, for example by the addition of HOBt or N-hydroxysuccinimide.

The starting materials of the formulae II and III are for the most part known. Insofar as they are not known, they can be prepared by known methods, for example the above-mentioned methods of peptide synthesis and of splitting protective groups.

If desired, a functionally modified amino and/or hydroxyl group in a compound of the formula I can be liberated

by solvolysis or hydrogenolysis in accordance with one of the methods described above.

Thus it is possible, in particular, to convert a compound of the formula I wherein X is other than H into a compound of the formula I ($X = H$), preferably by hydrogenolysis, if X is CBZ, otherwise by selective solvolysis. If X is BOC, the BOC group can be split off, for example by means of HCl in dioxane at room temperature.

Furthermore, it is possible, for example, to reduce keto compounds of the formula I ($R^4 = O$) to compounds of the formula I ($R^4 = H, OH$), for example by means of a complex metal hydride, such as $NaBH_4$, which does not simultaneously reduce the peptide carbonyl groups, in an inert solvent such as methanol at temperatures between about -10 and $+30^\circ$.

Keto compounds of the formula I ($R^4 = O$) can also be converted into compounds of the formula I ($R^4 = H, NH_2$) by reductive amination. Reductive amination can be carried out in a single stage or in several stages. Thus it is possible, for example, to treat the keto compound with ammonium salts, for example ammonium acetate, and $NaCNBH_3$, preferably in an inert solvent, for example an alcohol such as methanol, at temperatures between about 0 and 50° , in particular between 15 and 30° . It is also possible first to convert the keto compound into the oxime in a customary manner by means of hydroxylamine, and to reduce this oxime to the amine, for example by catalytic hydrogenation over Raney nickel.

It is also possible to convert a radical D into another radical D by treatment with esterifying, solvolysing or reducing agents. Thus an acid can be esterified, for example by means of an alcohol of the formula $A-OH$ or a diazoalkane, for example diazomethane, or an ester can be saponified to give the corresponding acid, for example by means of sodium hydroxide in aqueous dioxane solution at room temperature. It is also possible, for example, to convert a radical $R^9 = NH_2$ into a radical $R^9 = H$ by treatment with reducing agents, preferably with Raney nickel in an

alcohol such as isopropanol at temperatures between 20 and 120°.

A base of the formula I can be converted into the appropriate acid addition salt by means of an acid. Acids suitable for this reaction are, in particular, those which afford physiologically acceptable salts. Thus it is possible to use inorganic acids, for example sulfuric acid, nitric acid, hydrogen halide acids, such as hydrochloric or hydrobromic acid, phosphoric acids, such as orthophosphoric acid, or sulfamic acid, and also organic acids, in particular aliphatic, alicyclic, araliphatic, aromatic or heterocyclic monobasic or polybasic carboxylic, sulfonic or sulfuric acids, for example formic acid, acetic acid, propionic acid, pivalic acid, diethylacetic acid, malonic acid, succinic acid, pimelic acid, fumaric acid, maleic acid, lactic acid, tartaric acid, malic acid, benzoic acid, salicylic acid, 2-phenylpropionic acid, 3-phenylpropionic acid, citric acid, gluconic acid, ascorbic acid, nicotinic acid, isonicotinic acid, methanesulfonic acid, ethanesulfonic acid, ethanedisulfonic acid, 2-hydroxyethanesulfonic acid, benzenesulfonic acid, p-toluenesulfonic acid, naphthalene-monosulfonic and naphthalenedisulfonic acids or laurylsulfuric acid. Salts with physiologically unacceptable acids, for example picrates, can be used to isolate and/or purify the compounds of the formula I.

The new compounds of the formula I and their physiologically acceptable salts can be used for the preparation of pharmaceutical formulations by bringing them into a suitable dosage form together with at least one excipient or auxiliary and, if desired, together with one or more further active compound(s). The formulations thus obtained can be employed as medicaments in human or veterinary medicine. Suitable excipients are organic or inorganic substances which are suitable for enteral (for example oral or rectal) administration or parenteral administration or for administration in the form of an inhalation spray, and which do not react with the new compounds, for example water, vegetable oils, benzyl alcohols, polyethylene gly-

cols, glycerol triacetate and other fatty acid glycerides, gelatine, soya lecithin, carbohydrates, such as lactose or starch, magnesium stearate, talc or cellulose. Tablets, coated tablets, capsules, syrups, elixirs or drops are especially used for oral administration; lacquered tablets and capsules having coatings or capsule casings resistant to gastric juices are of particular interest. Suppositories are used for rectal administration; solutions, preferably oily or aqueous solutions, and also suspensions, emulsions or implants are used for parenteral application. Sprays containing the active compound either dissolved or suspended in a propellant gas mixture (for example fluorochlorohydrocarbons) can be used for administration as an inhalation spray. It is preferable in this regard to use the active compound in a micronized form, and one or more additional physiologically tolerable solvents can be present, for example ethanol. Inhalation solutions can be administered by means of customary inhalers. The new compounds can also be lyophilized and the resulting lyophilisates can be used, for example, for the preparation of injection formulations. The formulations indicated can be sterilized and/or can contain auxiliaries, such as preservatives, stabilizers and/or wetting agents, emulsifiers, salts for influencing the osmotic pressure, buffer substances, colorants and/or aroma substances. If desired, they can also contain one or more further active compounds, for example one or more vitamins.

As a rule, the substances according to the invention are administered analogously to other known, commercially available peptides, but, in particular, analogously to the compounds described in EP-A-77,028, preferably in dosages between about 100 mg and 30 g, especially between 500 mg and 5 g, per dosage unit. The daily dosage is preferably between about 2 and 600 mg/kg of body weight. The particular dose for each specific patient depends, however, on a very wide variety of factors, for example on the effectiveness of the particular compound employed, on the age, body weight, general state of health, sex, diet, time and means

of administration, excretion rate, combination of medications and the severity of the particular disease to which the therapy applies. Parenteral administration is preferred.

5 In the preceding and following text all temperatures are quoted in °C. In the following examples "customary working up" means as follows: if necessary, water is added, the mixture is neutralized and extracted with ether or methylene chloride, the phases are separated, the organic
10 phase is dried over sodium sulfate, filtered and evaporated, and the residue is purified by chromatography over silica gel and/or crystallization. $[\alpha] = [\alpha]_D^{20}$ in methanol, $c = 1$.

Example 1

15 The pH of a mixture of 978 mg of 2-[1S-(3S-hydroxy-4S-(N-tert.-butoxycarbonyl-L-phenylalanyl-N(imi)-(2,4-dinitrophenyl)-L-histidylamino)-5-cyclohexylpentanoylamino)-3-methylbutyl]-3H-quinazolin-4-one ["2-[1S-(BOC-Phe-imi-DNP-His-AHCP-amino)-3-methylbutyl]-3H-quinazolin-4-one";
20 obtainable by reacting BOC-Leu-OH with methyl anthranilate to give methyl 2-(BOC-Leu-amino)-benzoate (oil), reacting the latter with hydrazine hydrate to give 2-(1S-BOC-amino-3-methylbutyl)-3-amino-3H-quinazolin-4-one (m.p. 110-115° (decomp.); $[\alpha] -47.1^\circ$), boiling the latter for 5 hours
25 with Raney Ni in isopropanol with the formation of 2-(1S-BOC-amino-3-methylbutyl)-3H-quinazolin-4-one (m.p. 215° (decomp.); $[\alpha] -48.5^\circ$), splitting off the BOC group by means of 4N HCl in dioxane to give 2-(1S-amino-3-methylbutyl)-3H-quinazolin-4-one (dihydrochloride, m.p. 275° (decomp.);
30 $[\alpha] -28.4^\circ$), reacting the latter with BOC-AHCP-OH/DCCI/HOBt to give 2-(1S-BOC-AHCP-amino-3-methylbutyl)-3H-quinazolin-4-one, splitting off the BOC group and subjecting the product to a condensation reaction with BOC-imi-DNP-His-OH to give 2-(1S-BOC-imi-DNP-His-AHCP-amino-3-methylbutyl)-
35 3H-quinazolin-4-one, splitting off the BOC group again and reacting the product with BOC-Phe-OH], 2 g of 2-mercapto-ethanol, 20 ml of DMF and 20 ml of water is adjusted to 8 by stirring with aqueous Na₂CO₃ solution at 20°, and the

mixture is stirred for 2 hours at 20°. Working up in the customary manner gives 2-[1S-(3S-hydroxy-4S-(N-tert.-butoxycarbonyl-L-phenylalanyl-L-histidylamino)-5-cyclohexylpentanoylamino)-3-methylbutyl]-3H-quinazolin-4-one
5 [2-[1S-(BOC-Phe-His-AHCP-amino)-3-methylbutyl]-3H-quinazolin-4-one"], m.p. 147-149°.

The following are obtained analogously by cleaving the corresponding imi-DNP derivatives:

3-(BOC-Phe-His-AHCP-Ile-amino)-propane-1,2-diol, m.p. 180-
10 182°

3-(BOC-Phe-His-AHCP-Leu-amino)-propane-1,2-diol

2-(BOC-Phe-His-AHCP-Ile-amino)-ethanesulfonamide, m.p. 182°

2-(BOC-Phe-His-AHCP-Leu-amino)-ethanesulfonamide

o-(BOC-Phe-His-AHCP-Ile-amino)-benzenesulfonamide

15 o-(BOC-Phe-His-AHCP-Leu-amino)-benzenesulfonamide

m-(BOC-Phe-His-AHCP-Ile-amino)-benzenesulfonamide

m-(BOC-Phe-His-AHCP-Leu-amino)-benzenesulfonamide

N⁴-(BOC-Phe-His-AHCP-Ile)-sulfanilamide

20 N⁴-(BOC-Phe-His-AHCP-Leu)-sulfanilamide, m.p. 146-147°

p-(BOC-Phe-His-AHCP-Ile-aminomethyl)-benzenesulfonamide,
m.p. 227°

p-[2-(BOC-Phe-His-AHCP-Ile-amino)-ethyl]-benzenesulfonamide,

o-(BOC-Phe-His-AHCP-Ile-amino)-benzenesulfonic acid N-

25 methylamide

o-(BOC-Phe-His-AHCP-Leu-amino)-benzenesulfonic acid N-
methylamide

m-(BOC-Phe-His-AHCP-Ile-amino)-benzenesulfonic acid N-
methylamide

30 m-(BOC-Phe-His-AHCP-Leu-amino)-benzenesulfonic acid N-
methylamide

p-(BOC-Phe-His-AHCP-Ile-amino)-benzenesulfonic acid N-
methylamide, m.p. 157°

p-(BOC-Phe-His-AHCP-Leu-amino)-benzenesulfonic acid N-
35 methylamide

o-(BOC-Phe-His-AHCP-Ile-amino)-benzenesulfonic acid N,N-
dimethylamide

o-(BOC-Phe-His-AHCP-Leu-amino)-benzenesulfonic acid N,N-

dimethylamide

m-(BOC-Phe-His-AHCP-Ile-amino)-benzenesulfonic acid N,N-dimethylamide

m-(BOC-Phe-His-AHCP-Leu-amino)-benzenesulfonic acid N,N-dimethylamide

5

p-(BOC-Phe-His-AHCP-Ile-amino)-benzenesulfonic acid N,N-dimethylamide, m.p. 168°

p-(BOC-Phe-His-AHCP-Leu-amino)-benzenesulfonic acid N,N-dimethylamide

10

o-(morpholinoacetyl-Phe-His-AHCP-Ile-amino)-benzenesulfonamide

o-(morpholinoacetyl-Phe-His-AHCP-Leu-amino)-benzenesulfonamide

15

m-(morpholinoacetyl-Phe-His-AHCP-Ile-amino)-benzenesulfonamide

m-(morpholinoacetyl-Phe-His-AHCP-Leu-amino)-benzenesulfonamide

p-(morpholinoacetyl-Phe-His-AHCP-Ile-amino)-benzenesulfonamide

20

p-(morpholinoacetyl-Phe-His-AHCP-Leu-amino)-benzenesulfonamide

o-(morpholinoacetyl-Phe-His-AHCP-Ile-amino)-benzenesulfonic acid N-methylamide

25

o-(morpholinoacetyl-Phe-His-AHCP-Leu-amino)-benzenesulfonic acid N-methylamide

m-(morpholinoacetyl-Phe-His-AHCP-Ile-amino)-benzenesulfonic acid N-methylamide

m-(morpholinoacetyl-Phe-His-AHCP-Leu-amino)-benzenesulfonic acid N-methylamide

30

p-(morpholinoacetyl-Phe-His-AHCP-Ile-amino)-benzenesulfonic acid N-methylamide

p-(morpholinoacetyl-Phe-His-AHCP-Leu-amino)-benzenesulfonic acid N-methylamide

35

o-(morpholinoacetyl-Phe-His-AHCP-Ile-amino)-benzenesulfonic acid N,N-dimethylamide

o-(morpholinoacetyl-Phe-His-AHCP-Leu-amino)-benzenesulfonic acid N,N-dimethylamide

- m-(morpholinoacetyl-Phe-His-AHCP-Ile-amino)-benzenesulfonic acid N,N-dimethylamide
 m-(morpholinoacetyl-Phe-His-AHCP-Leu-amino)-benzenesulfonic acid N,N-dimethylamide
 5 p-(morpholinoacetyl-Phe-His-AHCP-Ile-amino)-benzenesulfonic acid N,N-dimethylamide
 p-(morpholinoacetyl-Phe-His-AHCP-Leu-amino)-benzenesulfonic acid N,N-dimethylamide
- 10 5-(BOC-Phe-His-AHCP-Ile-amino)-furan-2-sulfonamide
 5-(BOC-Phe-His-AHCP-Leu-amino)-furan-2-sulfonamide
 5-(BOC-Phe-His-AHCP-Ile-amino)-thiophene-2-sulfonamide
 5-(BOC-Phe-His-AHCP-Leu-amino)-thiophene-2-sulfonamide
 5-(BOC-Phe-His-AHCP-Ile-amino)-thiophene-3-sulfonamide
 15 5-(BOC-Phe-His-AHCP-Leu-amino)-thiophene-3-sulfonamide
 2-(BOC-Phe-His-AHCP-Ile-amino)-pyridine-5-sulfonamide
 2-(BOC-Phe-His-AHCP-Leu-amino)-pyridine-5-sulfonamide
 p-(BOC-Phe-His-AHCP-Ile-amino)-benzamide, m.p. 228°
 p-(BOC-Phe-His-AHCP-Leu-amino)-benzamide
- 20 p-(BOC-Phe-His-AHCP-Ile-amino)-benzenephosphonic acid diamide
 p-(BOC-Phe-His-AHCP-Leu-amino)-benzenephosphonic acid diamide
- 25 2-[1S-(BOC-Phe-His-AHCP-amino)-ethyl]-3H-quinazolin-4-one
 2-[1S-(BOC-Phe-His-AHCP-amino)-ethyl]-3-amino-3H-quinazolin-4-one [obtainable via methyl 2-(BOC-Ala-amino)-benzoate (m.p. 110-112°)]
 2-[1S-(BOC-Phe-His-AHCP-amino)-2-methylpropyl]-3H-quinazolin-4-one
 30 2-[1S-(BOC-Phe-His-AHCP-amino)-2-methylpropyl]-3-amino-3H-quinazolin-4-one, m.p. 127° [obtainable via methyl 2-(BOC-Val-amino)-benzoate (m.p. 151-155°)]
 2-[1S-(BOC-Phe-His-AHCP-amino)-2S-methylbutyl]-3H-quinazolin-4-one
 35 2-[1S-(BOC-Phe-His-AHCP-amino)-2S-methylbutyl]-3-amino-3H-quinazolin-4-one, m.p. 120° [decomp.; obtainable via 2-(1S-BOC-amino-2S-methylbutyl)-3-amino-3H-quinazolin-4-one

(m.p. 110-115°; [α] -41.9°) and 2-(1S-amino-2S-methylbutyl)-3-amino-3H-quinazolin-4-one (m.p. 105° (decomp.); [α] -4.4°)

2-[1S-(BOC-Phe-His-AHCP-amino)-3-methylbutyl]-3-amino-3H-
5 quinazolin-4-one, m.p. 125-128°

2-[1S-(BOC-Phe-His-AHCP-amino)-2-phenylethyl]-3H-quinazolin-4-one

2-[1S-(BOC-Phe-His-AHCP-amino)-2-phenylethyl]-3-amino-3H-
10 quinazolin-4-one, m.p. 198° [obtainable via methyl 2-(BOC-Phe-amino)-benzoate (m.p. 145-147°)]

2-[1S-(BOC-Phe-His-AHCP-amino)-2S-methylbutyl]-6-sulfamoyl-7-chloro-3H-quinazolin-4-one

2-[1S-(BOC-Phe-His-AHCP-amino)-2S-methylbutyl]-3-amino-6-
15 sulfamoyl-7-chloro-3H-quinazolin-4-one

2-[1S-(BOC-Phe-His-AHCP-amino)-3-methylbutyl]-6-sulfamoyl-7-chloro-3H-quinazolin-4-one

2-[1S-(BOC-Phe-His-AHCP-amino)-3-methylbutyl]-3-amino-6-sulfamoyl-7-chloro-3H-quinazolin-4-one

20

2-[1S-(N-(2-pyrrolidino-3-phenyl-propionyl)-His-AHCP-amino)-2S-methylbutyl]-3H-quinazolin-4-one

2-[1S-(N-(2-pyrrolidino-3-phenyl-propionyl)-His-AHCP-amino)-2S-methylbutyl]-3-amino-3H-quinazolin-4-one

25 2-[1S-(N-(2-pyrrolidino-3-phenyl-propionyl)-His-AHCP-amino)-3-methylbutyl]-3H-quinazolin-4-one

2-[1S-(N-(2-pyrrolidino-3-phenyl-propionyl)-His-AHCP-amino)-3-methylbutyl]-3-amino-3H-quinazolin-4-one

2-[1S-(N-(2-piperidino-3-phenyl-propionyl)-His-AHCP-amino)-2S-methylbutyl]-3H-quinazolin-4-one

2-[1S-(N-(2-piperidino-3-phenyl-propionyl)-His-AHCP-amino)-2S-methylbutyl]-3-amino-3H-quinazolin-4-one

2-[1S-(N-(2-piperidino-3-phenyl-propionyl)-His-AHCP-amino)-3-methylbutyl]-3H-quinazolin-4-one

35 2-[1S-(N-(2-piperidino-3-phenyl-propionyl)-His-AHCP-amino)-3-methylbutyl]-3-amino-3H-quinazolin-4-one

2-[1S-(N-(2-morpholino-3-phenyl-propionyl)-His-AHCP-amino)-2S-methylbutyl]-3H-quinazolin-4-one

- 2-[1S-(N-(2-morpholino-3-phenyl-propionyl)-His-AHCP-amino)-
2S-methylbutyl]-3-amino-3H-quinazolin-4-one
- 2-[1S-(N-(2-morpholino-3-phenyl-propionyl)-His-AHCP-amino)-
3-methylbutyl]-3H-quinazolin-4-one
- 5 2-[1S-(N-(2-morpholino-3-phenyl-propionyl)-His-AHCP-amino)-
3-methylbutyl]-3-amino-3H-quinazolin-4-one
- 2-[1S-(N-(2-benzyl-hexanoyl)-His-AHCP-amino)-2S-methyl-
butyl]-3H-quinazolin-4-one
- 2-[1S-(N-(2-benzyl-hexanoyl)-His-AHCP-amino)-2S-methyl-
10 butyl]-3-amino-3H-quinazolin-4-one
- 2-[1S-(N-(2-benzyl-hexanoyl)-His-AHCP-amino)-3-methyl-
butyl]-3H-quinazolin-4-one
- 2-[1S-(N-(2-benzyl-hexanoyl)-His-AHCP-amino)-3-methyl-
butyl]-3-amino-3H-quinazolin-4-one
- 15 2-[1S-(N-(2-butoxy-3-phenyl-propionyl)-His-AHCP-amino)-2S-
methylbutyl]-3H-quinazolin-4-one
- 2-[1S-(N-(2-butoxy-3-phenyl-propionyl)-His-AHCP-amino)-2S-
methylbutyl]-3-amino-3H-quinazolin-4-one
- 2-[1S-(N-(2-butoxy-3-phenyl-propionyl)-His-AHCP-amino)-
20 3-methylbutyl]-3H-quinazolin-4-one
- 2-[1S-(N-(2-butoxy-3-phenyl-propionyl)-His-AHCP-amino)-
3-methylbutyl]-3-amino-3H-quinazolin-4-one
- 2-[1S-(N-(2-butylthio-3-phenyl-propionyl)-His-AHCP-amino)-
2S-methylbutyl]-3H-quinazolin-4-one
- 25 2-[1S-(N-(2-butylthio-3-phenyl-propionyl)-His-AHCP-amino)-
2S-methylbutyl]-3-amino-3H-quinazolin-4-one
- 2-[1S-(N-(2-butylthio-3-phenyl-propionyl)-His-AHCP-amino)-
3-methylbutyl]-3H-quinazolin-4-one
- 2-[1S-(N-(2-butylthio-3-phenyl-propionyl)-His-AHCP-amino)-
30 3-methylbutyl]-3-amino-3H-quinazolin-4-one

Example 2

- 1 g of 2-[1S-(BOC-Phe-(imi-BOM-His)-AHCP-amino)-2S-
methylbutyl]-3-amino-3H-quinazoline-4-one [m.p. 146°;
[α] -37.2°; obtainable from 2-(1S-amino-2S-methylbutyl)-
35 3-amino-3H-quinazolin-4-one and BOC-Phe-(imi-BOM-His)-AHCP-
OH] is dissolved in 15 ml of methanol, and the mixture is
hydrogenated over 0.5 % Pd/C at 20° and 1 bar until absorp-
tion ceases, and is filtered and

(BOC-Phe-His-AHCP-amino)-2S-methylbutyl]-3-amino-3H-quinazolin-4-one, m.p. 120° (decomp.).

The other compounds indicated in Example 1 are obtained analogously by cleaving the corresponding imi-BOH derivatives, as are also the following:

p-(BOC-Phe-His-AHCP-Ile-amino)-benzenesulfonanilide

p-(BOC-Phe-His-AHCP-Ile-amino)-benzenesulfonic acid o-sulfamoylanilide

10 p-(BOC-Phe-His-AHCP-Ile-amino)-benzenesulfonic acid p-sulfamoylanilide

p-(BOC-Phe-His-AHCP-Ile-amino)-benzenesulfonic acid N-(2-thienyl)-amide

15 p-(BOC-Phe-His-AHCP-Ile-amino)-benzenesulfonic acid N-(2-thiazolyl)-amide

p-(BOC-Phe-His-AHCP-Ile-amino)-benzenesulfonic acid N-(3-isoxazolyl)-amide

p-(BOC-Phe-His-AHCP-Ile-amino)-benzenesulfonic acid N-(5-methyl-3-isoxazolyl)-amide

20 p-(BOC-Phe-His-AHCP-Ile-amino)-benzenesulfonic acid N-(3,4-dimethyl-5-isoxazolyl)-amide

p-(BOC-Phe-His-AHCP-Ile-amino)-benzenesulfonic acid N-(4-pyridyl)-amide

25 p-(BOC-Phe-His-AHCP-Ile-amino)-benzenesulfonic acid N-(2-pyrimidinyl)-amide, m.p. 192°

p-(BOC-Phe-His-AHCP-Ile-amino)-benzenesulfonic acid N-(4-methyl-2-pyrimidinyl)-amide

p-(BOC-Phe-His-AHCP-Ile-amino)-benzenesulfonic acid N-(4,6-dimethyl-2-pyrimidinyl)-amide

30 p-(BOC-Phe-His-AHCP-Ile-amino)-benzenesulfonic acid N-(2,6-dimethyl-4-pyrimidinyl)-amide.

Example 3

1.01 g of N-methylmorpholine is added to a solution, in 60 ml of CH₂Cl₂, of 4.58 g of 3-(H-Gly-AHCP-Ile-amino)-propane-1,2-diol [obtainable by reacting BOC-Gly-AHCP-OH with 3-(H-Ile-amino)-propane-1,2-diol to give 3-(BOC-Gly-AHCP-Ile-amino)-propane-1,2-diol and splitting off the BOC group]. 2.65 g of BOC-Phe-OH, 1.35 g of HOBT and

a solution of 2.06 g of DCCI in 50 ml of CH₂Cl₂ are added with stirring, the mixture is stirred for 14 hours at 4°, the precipitated dicyclohexylurea is filtered off and the filtrate is evaporated. Customary working up gives 3-(BOC-Phe-Gly-AHCP-Ile-amino)-propane-1,2-diol, m.p. 104-106°.

Example 4

N⁴-(BOC-Phe-Gly-AHCP-Leu)-sulfanilamide, m.p. 142-144°, is obtained analogously to Example 3 from BOC-Phe-Gly-OH and N⁴-(H-AHCP-Leu)-sulfanilamide.

10 The following are obtained analogously:

N⁴-(BOC-Phe-Gly-AHCP-Ile)-sulfanilamide

N⁴-(POA-Phe-Abu-AHCP-Leu)-sulfanilamide

N⁴-(ETOC-Phe-Ada-AHCP-Leu)-sulfanilamide

N⁴-(IPOC-Phe-Ala-AHCP-Leu)-sulfanilamide

15 N⁴-(CBZ-Phe-Cal-AHCP-Leu)-sulfanilamide

N⁴-(acetyl-Phe-(N-is-methyl-His)-AHCP-Leu)-sulfanilamide

N⁴-(ETNC-Phe-Ile-AHCP-Leu)-sulfanilamide

N⁴-(IPNC-Phe-Leu-AHCP-Leu)-sulfanilamide

N⁴-(MC-Phe-tert.-Leu-AHCP-Leu)-sulfanilamide

20 N⁴-(PBB-Phe-Met-AHCP-Leu)-sulfanilamide

N⁴-(4-phenylbutyryl-Phe-αNal-AHCP-Leu)-sulfanilamide

N⁴-(2-benzyl-3-phenylpropionyl-Phe-βNal-AHCP-Leu)-sulfanilamide

N⁴-(morpholinoacetyl-Phe-βAla-AHCP-Leu)-sulfanilamide, m.p. 192°

25 N⁴-[2-(2-phenylethyl)-4-phenylbutyryl-Phe-Nbg-AHCP-Leu]-sulfanilamide

N⁴-[2-(2-naphthylmethyl)-4-phenylbutyryl-Phe-Nle-AHCP-Leu]-sulfanilamide

N⁴-[propionyl-Phe-(N-Me-His)-AHCP-Leu]-sulfanilamide

30 N⁴-[butyryl-Phe-(N-Me-Phe)-AHCP-Leu]-sulfanilamide

N⁴-(isobutyryl-Phe-Phe-AHCP-Leu)-sulfanilamide

N⁴-(cyclopentylcarbonyl-Phe-Pro-AHCP-Leu)-sulfanilamide

N⁴-(cyclohexylcarbonyl-Phe-Ser-AHCP-Leu)-sulfanilamide

N⁴-(benzoyl-Phe-Thr-AHCP-Leu)-sulfanilamide

35 N⁴-(phenylacetyl-Phe-Tic-AHCP-Leu)-sulfanilamide

N⁴-(2-phenylpropionyl-Phe-Trp-AHCP-Leu)-sulfanilamide

N⁴-(3-phenylpropionyl-Phe-Tyr-AHCP-Leu)-sulfanilamide

N⁴-(2-p-fluorophenylpropionyl-Phe-Val-AHCP-Leu)-sulfanil-

Example 5

N^4 -(BOC-Phe-Gly-AHCP-Leu)-sulfanilamide, m.p. 142-144°, is obtained analogously to Example 3 from BOC-Phe-Gly-AHCP-OH and N^4 -(H-Leu)-sulfanilamide.

5 The following are obtained analogously:

N^4 -(BOC-Phe-BAla-AHCP-Leu)-sulfanilamide, m.p. 196°

N^4 -(BOC-Phe-Gly-AHCP-Ile)-sulfanilamide

5-(BOC-Phe-Gly-AHCP-Ile-amino)-thiophene-3-sulfonamide

2-[1S-(BOC-Phe-Gly-AHCP-amino)-2-methylpropyl]-3H-quinazolin-4-one

2-[1S-(BOC-Phe-Gly-AHCP-amino)-2-methylpropyl]-3-amino-3H-quinazolin-4-one

2-[1S-(BOC-Phe-Gly-AHCP-amino)-2S-methylbutyl]-3H-quinazolin-4-one

15 2-[1S-(BOC-Phe-Gly-AHCP-amino)-2S-methylbutyl]-3-amino-3H-quinazolin-4-one, m.p. 115° (decomp.)

2-[1S-(BOC-Phe-Gly-AHCP-amino)-3-methylbutyl]-3H-quinazolin-4-one

2-[1S-(BOC-Phe-Gly-AHCP-amino)-3-methylbutyl]-3-amino-3H-quinazolin-4-one

2-[1S-(BOC-Phe-Gly-AHCP-amino)-2-phenylethyl]-3H-quinazolin-4-one

2-[1S-(BOC-Phe-Gly-AHCP-amino)-2-phenylethyl]-3-amino-3H-quinazolin-4-one

2-[1S-(BOC-Phe-Gly-AHCP-amino)-2S-methylbutyl]-6-sulfamoyl-7-chloro-3H-quinazolin-4-one

2-[1S-(BOC-Phe-Gly-AHCP-amino)-2S-methylbutyl]-3-amino-6-sulfamoyl-7-chloro-3H-quinazolin-4-one

30 2-[1S-(BOC-Phe-Gly-AHCP-amino)-3-methylbutyl]-6-sulfamoyl-7-chloro-3H-quinazolin-4-one

2-[1S-(BOC-Phe-Gly-AHCP-amino)-3-methylbutyl]-3-amino-6-sulfamoyl-7-chloro-3H-quinazolin-4-one

2-(BOC-Phe-Gly-AHCP-Ile-aminomethyl)-3H-quinazolin-4-one

35 2-(BOC-Phe-Gly-AHCP-Ile-aminomethyl)-3-amino-3H-quinazolin-4-one, m.p. 188°

2-[1S-(N-(2-pyrrolidino-3-phenyl-propionyl)-Gly-AHCP-amino)-2S-methylbutyl]-3H-quinazolin-4-one

2-[1S-(N-(2-pyrrolidino-3-phenyl-propionyl)-Gly-AHCP-amino)-

- 2-[1S-(N-(2-pyrrolidino-3-phenyl-propionyl)-Gly-AHCP-amino)-
3-methylbutyl]-3H-quinazolin-4-one
- 2-[1S-(N-(2-pyrrolidino-3-phenyl-propionyl)-Gly-AHCP-amino)-
3-methylbutyl]-3-amino-3H-quinazolin-4-one
- 5 2-[1S-(N-(2-piperidino-3-phenyl-propionyl)-Gly-AHCP-amino)-
2S-methylbutyl]-3H-quinazolin-4-one
- 2-[1S-(N-(2-piperidino-3-phenyl-propionyl)-Gly-AHCP-amino)-
2S-methylbutyl]-3-amino-3H-quinazolin-4-one
- 2-[1S-(N-(2-piperidino-3-phenyl-propionyl)-Gly-AHCP-amino)-
10 3-methylbutyl]-3H-quinazolin-4-one
- 2-[1S-(N-(2-piperidino-3-phenyl-propionyl)-Gly-AHCP-amino)-
3-methylbutyl]-3-amino-3H-quinazolin-4-one
- 2-[1S-(N-(2-morpholino-3-phenyl-propionyl)-Gly-AHCP-amino)-
2S-methylbutyl]-3H-quinazolin-4-one
- 15 2-[1S-(N-(2-morpholino-3-phenyl-propionyl)-Gly-AHCP-amino)-
2S-methylbutyl]-3-amino-3H-quinazolin-4-one
- 2-[1S-(N-(2-morpholino-3-phenyl-propionyl)-Gly-AHCP-amino)-
3-methylbutyl]-3H-quinazolin-4-one
- 2-[1S-(N-(2-morpholino-3-phenyl-propionyl)-Gly-AHCP-amino)-
20 3-methylbutyl]-3-amino-3H-quinazolin-4-one
- 2-[1S-(N-(2-benzyl-hexanoyl)-Gly-AHCP-amino)-2S-methyl-
butyl]-3H-quinazolin-4-one
- 2-[1S-(N-(2-benzyl-hexanoyl)-Gly-AHCP-amino)-2S-methyl-
25 butyl]-3-amino-3H-quinazolin-4-one
- 2-[1S-(N-(2-benzyl-hexanoyl)-Gly-AHCP-amino)-3-methyl-
butyl]-3H-quinazolin-4-one
- 2-[1S-(N-(2-benzyl-hexanoyl)-Gly-AHCP-amino)-3-methyl-
butyl]-3-amino-3H-quinazolin-4-one
- 30 2-[1S-(N-(2-butoxy-3-phenyl-propionyl)-Gly-AHCP-amino)-
2S-methylbutyl]-3H-quinazolin-4-one
- 2-[1S-(N-(2-butoxy-3-phenyl-propionyl)-Gly-AHCP-amino)-
2S-methylbutyl]-3-amino-3H-quinazolin-4-one
- 2-[1S-(N-(2-butoxy-3-phenyl-propionyl)-Gly-AHCP-amino)-
35 3-methylbutyl]-3H-quinazolin-4-one
- 2-[1S-(N-(2-butoxy-3-phenyl-propionyl)-Gly-AHCP-amino)-
3-methylbutyl]-3-amino-3H-quinazolin-4-one
- 2-[1S-(N-(2-butylthio-3-phenyl-propionyl)-Gly-AHCP-amino)-

2S-methylbutyl]-3H-quinazolin-4-one

2-[1S-(N-(2-butylthio-3-phenyl-propionyl)-Gly-AHCP-amino)-
2S-methylbutyl]-3-amino-3H-quinazolin-4-one

5 2-[1S-(N-(2-butylthio-3-phenyl-propionyl)-Gly-AHCP-amino)-
3-methylbutyl]-3H-quinazolin-4-one

2-[1S-(N-(2-butylthio-3-phenyl-propionyl)-Gly-AHCP-amino)-
3-methylbutyl]-3-amino-3H-quinazolin-4-one

Example 6

10 3-(BOC-Phe-Gly-AHCP-Ile-Met-amino)-propane-1,2-diol
is obtained analogously to Example 3 from BOC-Phe-Gly-AHCP-
Ile-OH and 3-(H-Met-amino)-propane-1,2-diol.

Example 7

15 3-(BOC-Phe-Gly-AHCP-Ile-amino)-propane-1,2-diol,
m.p. 104-106°, is obtained analogously to Example 3 from
BOC-Phe-Gly-AHCP-Ile-OH and 3-aminopropane-1,2-diol.

The following are obtained analogously:

3-(BOC-Phe-Gly-AHCP-Abu-amino)-propane-1,2-diol
3-(BOC-Phe-Gly-AHCP-Ala-amino)-propane-1,2-diol
3-(BOC-Phe-Gly-AHCP-Cal-amino)-propane-1,2-diol
20 3-(BOC-Phe-Gly-AHCP-His-amino)-propane-1,2-diol
3-(BOC-Phe-Gly-AHCP-Leu-amino)-propane-1,2-diol
3-(BOC-Phe-Gly-AHCP-Met-amino)-propane-1,2-diol
3-(BOC-Phe-Gly-AHCP-Nle-amino)-propane-1,2-diol
3-(BOC-Phe-Gly-AHCP-Phe-amino)-propane-1,2-diol
25 3-(BOC-Phe-Gly-AHCP-Trp-amino)-propane-1,2-diol
3-(BOC-Phe-Gly-AHCP-Tyr-amino)-propane-1,2-diol
3-(BOC-Phe-Gly-AHCP-Val-amino)-propane-1,2-diol
3-(BOC-Phe-Gly-AHCH-Ile-amino)-propane-1,2-diol
3-(BOC-Phe-Gly-Sta-Ile-amino)-propane-1,2-diol
30 3-(BOC-Phe-Gly-AHPP-Ile-amino)-propane-1,2-diol.

Example 8

35 A solution of 1 g of 2-[1S-(BOC-Phe-His-AHCP-amino)-
3-methylbutyl]-3H-quinazolin-4-one in 20 ml of 4N HCl in
dioxane is stirred for 30 minutes at 20° and is then eva-
porated. This gives 2-[1S-(H-Phe-His-AHCP-amino)-3-methyl-
butyl]-3H-quinazolin-4-one.

The following are obtained analogously by cleaving
the corresponding BOC derivatives.

3-(H-Phe-Gly-AHCP-Ile-amino)-propane-1,2-diol

3-(H-Phe-His-AHCP-Ile-amino)-propane-1,2-diol

N⁴-(H-Phe-Gly-AHCP-Leu)-sulfanilamide

N⁴-(H-Phe-His-AHCP-Leu)-sulfanilamide

5

2-[1S-(H-Phe-His-AHCP-amino)-3-methylbutyl]-3-amino-3H-quinazolin-4-one

2-[1S-(H-Phe-His-AHCP-amino)-2S-methylbutyl]-3H-quinazolin-4-one

10 2-[1S-(H-Phe-His-AHCP-amino)-2S-methylbutyl]-3-amino-3H-quinazolin-4-one.

Example 9

1 g of 2-[1S-(CBZ-Phe-His-AHCP-amino)-3-methylbutyl]-3H-quinazolin-4-one is dissolved in 15 ml of ethanol,
15 and the mixture is hydrogenated over 0.5 g of 10% Pd/C at 20° and 1 bar for 3 hours and is filtered and evaporated to give, after purification by chromatography, 2-[1S-(H-Phe-His-AHCP-amino)-3-methylbutyl]-3H-quinazolin-4-one.

Example 10

20 70 mg of hydroxylamine hydrochloride are added to a solution of 826 mg of 2-[1S-(3-Oxo-4S-BOC-Phe-His-amino-5-cyclohexylpentanoylamino)-3-methylbutyl]-3H-quinazolin-4-one and 1.43 g of Na₂CO₃ · 10 H₂O in 5 ml of methanol and 5 ml of water, and the mixture is stirred for 14 hours at
25 20°. The precipitated oxime is filtered off, dried, dissolved in 10 ml of methanol and hydrogenated over 0.5 g of Raney Ni at 20° and 5 bar. The catalyst is filtered off, the filtrate is evaporated and the resulting mixture is separated over silica gel to give 2-[1S-(3S-amino-4S-BOC-Phe-His-amino-5-cyclohexylpentanoylamino)-3-methylbutyl]-
30 3H-quinazolin-4-one ["2-[1S-(BOC-Phe-His-DACP-amino)-3-methylbutyl]-3H-quinazolin-4-one"]; the 3R-amino epimer is also obtained.

35 The following are obtained analogously from the corresponding oxo compounds:

2-[1S-(BOC-Phe-His-DACH-amino)-3-methylbutyl]-3H-quinazolin-4-one

2-[1S-(BOC-Phe-His-DAMH-amino)-3-methylbutyl]-3H-quinazolin-4-one

2-[1S-(BOC-Phe-His-DAPP-amino)-3-methylbutyl]-3H-quinazolin-4-one.

5 Example 11

0.5 g of Raney nickel, moistened with 0.5 g of isopropanol, is added to a solution of 1 g of 2-[1S-(BOC-Phe-His-AHCP-amino)-3-methylbutyl]-3-amino-3H-quinazolin-4-one in 500 ml of isopropanol, and the mixture is boiled for 10 5 hours. After filtration, the mixture is evaporated and worked up in the customary manner to give 2-[1S-(BOC-Phe-His-AHCP-amino)-3-methylbutyl]-3H-quinazolin-4-one, m.p. 147-149°.

The examples below relate to pharmaceutical formulations. 15

Example A: Injection vials

The pH of a solution of 100 g of 2-[1S-(BOC-Phe-His-AHCP-amino)-3-methylbutyl]-3H-quinazolin-4-one and 5 g of disodium hydrogenphosphate in 4 l of twice distilled 20 water is adjusted to 6.5 with 2N hydrochloric acid, and the solution is filtered under sterile conditions and filled into injection vials. These are lyophilized under sterile conditions and closed in a sterile manner. Each injection vial contains 500 mg of active compound.

25 Example B: Suppositories

A mixture of 50 g of 2-[1S-(BOC-Phe-His-AHCP-amino)-3-methylbutyl]-3-amino-3H-quinazolin-4-one with 10 g of soya lecithin and 140 g of cocoa butter is melted, poured into moulds and allowed to cool. Each suppository contains 250 30 mg of active compound.

Merck Patent Gesellschaft
mit beschränkter Haftung
6100 Darmstadt


The claims defining the invention are as follows:

1. Amino acid derivatives of the formula I



wherein

- X is H , $R^1-O-C_{mH_{2m}}-CO-$, $R^1-C_{mH_{2m}}-O-CO-$,
 $R^1-C_{mH_{2m}}-CO-$, R^1-SO_2- or
 $(R^1-C_{mH_{2m}}-(T)_x-C_{rH_{2r}})-L(R^7-C_{pH_{2p}})-C_{tH_{2t}}-CO-$,
 Z is 0 to 4 amino acid radicals attached to one another by a peptide linkage and selected from the group consisting of Abu, Ada, Ala, BAla, Arg, Asn, Asp, Bia, Cal, Dab, Gln, Glu, Gly, His, N(im)-alkyl-His, Ile, Leu, tert.-Leu, Lys, Met, α Nal, BNal, Nbg, Nle, Orn, Phe, Pro, Ser, Thr, Tic, Trp, Tyr and Val,
 E is 0 to 2 amino acid radicals attached to one another by a peptide linkage and selected from the group consisting of Abu, Ala, Cal, His, Ile, Leu, Met, Nle, Phe, Trp, Tyr and Val,
 D is $-CH_2-CHOH-CH_2OH$, $-C_2H_2-SO_2-R^{14}$, a phenyl- C_6H_5- , furyl- C_4H_3- , thienyl- C_4H_3- or pyridyl- C_4H_3- radical which is substituted by one or two $R^{14}-SO_2-$ groups or an $R^{14}-CO-$ group or an $(R^{14})_2-PO-$ group and, if desired additionally by an Hal atom, or D is



R^1 , R^3 , R^7 and R^8 are each H, A, Ar, Ar-alkyl, Het,

Het-alkyl or cycloalkyl having 3-7 C atoms, cycloalkylalkyl having 4-11 C atoms, bicycloalkyl or tricycloalkyl having in each case 7-14 C atoms or bicycloalkylalkyl or tricycloalkylalkyl having in each case 8-18 C atoms, each of which is unsubstituted or monosubstituted or polysubstituted by A, AO and/or Hal,

R^2 , R^5 and R^6 are each H or A,

R^4 is (H, OH), (H, NH_2) or =O,

R^9 is H, NH_2 , NHA or NA_2 ,

R^{10} , R^{11} , R^{12} and R^{13} are each H, Hal, OH, OA, NH_2 , SH, SA, SO_2NH_2 , CF_3 , CN, COOH or COOA,

R^{14} is OH, OA, NH_2 , NHA, NA_2 , NHCycloalkyl having 3-7 C atoms, N(cycloalkyl)₂ having 6-14 C atoms, pyrrolidino, piperidino, hexahydroazepino, morpholino, thiomorpholino, piperazino, N-A-piperazino, NHA_{Ar} or NHHet,

L is CH or N,

T is O, S, NH or NA,

n is 1 or 2,

m, p, r and t are each 0, 1, 2, 3, 4 or 5,

x is 0 or 1,

y is 0, 1 or 2,

z is 2, 3, 4, 5 or 6,

Ar is phenyl which is unsubstituted or monosubstituted or polysubstituted by A, AO, Hal, CF_3 , OH, H_2NSO_2 and/or NH_2 or unsubstituted naphthyl,

Het is a saturated or unsaturated 5-membered or 6-membered heterocyclic radical which has 1-4 N, O and/or S atoms, which can be condensed with a benzene ring and/or can be monosubstituted or polysubstituted by A, AO, Hal, CF_3 , HO, O_2N , carbonyl oxygen, H_2N , HAN, A_2N , AcNH, AS, ASO, ASO_2 , HOOC, AOOC, CN, H_2NCO , H_2NSO_2 , ASO_2NH , Ar or Ar-alkenyl, hydroxyalkyl and/or aminoalkyl having in each case 1-8 C atoms, and/or in which the N and/or S heteroatoms can also be oxidized,

Hal is F, Cl, Br or I,

Ac is A-CO-, Ar-CO- or A-NH-CO-,
alkyl- is an alkylene group having 1-4 C atoms and
A is alkyl having 1-8 C atoms,
and wherein it is also possible for one or more -NA-CO-
groups to replace one or more -NH-CO- groups, and salts
thereof.

2. a) 2-[1S-(BOC-Phe-His-AHCP-amino)-2S-methyl-
butyl]-3H-quinazol in-4-one;
- b) 2-[1S-(BOC-Phe-His-AHCP-amino)-2S-methyl-
butyl]-3-amino-3H-quinazol in-4-one;
- c) 2-[1S-(BOC-Phe-His-AHCP-amino)-3-methyl-
butyl]-3H-quinazol in-4-one; or
- d) 2-[1S-(BOC-Phe-His-AHCP-amino)-3-methyl-
butyl]-3-amino-3H-quinazol in-4-one.

3. Process for the preparation of an amino acid derivative
of the formula I and salts thereof, characterized in that
this amino acid derivative is liberated from one of its
functional derivatives by treatment with a solvolysing or
hydrogenolysing agent, or in that a carboxylic acid of the

formula II



wherein G^1 is

(a) Z^1 ,

(b) Z,

(c) Z-W,


(d) Z-W-E¹,

(e) Z-W-E

II

and

W is



is reacted with an amino compound of the formula III



III

wherein G^2 is

(a) $\text{-Z}^2\text{-W-E-NR}^6\text{-D,}$

- (b) $-W-E-NR^6-D$,
- (c) $-E-NR^6-D$,
- (d) $-E^2-NR^6-D$,
- (e) $-NR^6-D$,

E^1 and E^2 are each one amino acid radical selected from the group consisting of Abu, Ala, Cal, His, Ile, Leu, Met, Nle, Phe, Trp, Tyr and Val in such a manner that $E^1 + E^2$ together are E,

Z^1 and Z^2 are each 1 to 3 amino acid radicals selected from the group consisting of Abu, Ada, Ala, SAla, Arg, Asn, Asp, Bia, Cal, Dab, Gln, Glu, Gly, His N(im)-alkyl-His, Ile, Leu, tert -Leu, Lys, Met, α Nal, SAla, Nbg, Nle, Orn, Phe, Pro, Ser, Thr, Tic, Trp, Tyr and Val in such a manner that $Z^1 + Z^2$ together are Z, and in that, if appropriate, a functionally modified amino and/or hydroxyl group in a compound of the formula I is Liberated by treatment with solvolysing or hydrogenolysing agents and/or, in order to prepare a compound of the formula I wherein $R^4 = (H, OH)$ or (H, NH_2) , an aminoketo acid derivative of the formula I wherein $R^4 = O$ is reduced or reductively aminated and/or a radical D is converted into an- other radical D by treatment with esterifying, solvolysing or reducing agents and/or a compound of the formula I is converted into one of its salts by treatment with an acid.

4. Process for the preparation of pharmaceutical formulations, characterized in that a compound of the formula I and/or one of its physiologically acceptable salts is brought into a suitable dosage form together with at least one solid, liquid or semi-liquid excipient or auxiliary and, if appropriate, in combination with one or more further active compound(s).

5. Pharmaceutical formulation characterized in that it contains at least one compound of the formula I and/or one of its physiologically acceptable salts.

6. The use of compounds of the formula I or of physiologically acceptable salts thereof for the preparation of a medicament.

7. The use of compounds of the formula I or of physiologically acceptable salts thereof in combating renin-dependent hypertension or hyperaldosteronism.

DATED this 1st day of July, 1991.

MERCK PATENT
GESELLSCHAFT MIT
BESCHRANKTER HUNFTUNG
By Its Patent Attorneys
ARTHUR S. CAVE & CO.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418
1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095
2096
2097
2098
2099
2100
2101
2102
2103
2104
2105
2106
2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2118
2119
2120
2121
2122
2123
2124
2125
2126
2127
2128
2129
2130
2131
2132
2133
2134
2135
2136
2137
2138
2139
2140
2141
2142
2143
2144
2145
2146
2147
2148
2149
2150
2151
2152
2153
2154
2155
2156
2157
2158
2159
2160
2161
2162
2163
2164
2165
2166
2167
2168
2169
2170
2171
2172
2173
2174
2175
2176
2177
2178
2179
2180
2181
2182
2183
2184
2185
2186
2187
2188
2189
2190
2191
2192
2193
2194
2195
2196
2197
2198
2199
2200
2201
2202
2203
2204
22

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☐ FADED TEXT OR DRAWING
- ☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☒ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☐ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.